

Enumerating maximal consistent closed sets in closure systems

AICoLoCo


Lhouari Nourine and *Simon Vilmin*.

LIMOS, UCA

May 2021

Implications and inconsistency

- ▶ A set X of attributes with implications Σ , and closure system \mathcal{F}
- ▶ A *consistency graph* G (over X)
- ▶ Problem MCCENUM : enumerate *maximal consistent closed sets* of \mathcal{F} w.r.t G



Problem - Maximal consistent closed sets enumeration (MCCENUM)

- ▶ **Input:** A set of implications Σ over X , a consistency graph $G = (X, E)$.
- ▶ **Output:** maximal consistent closed sets of \mathcal{F} w.r.t G , denoted $\max\text{CC}(\Sigma, G)$.


Origins:

- ▶ Representation for median-semilattices [Barthélemy, Constantin, 1993], [Nielsen et al., 1981]
- ▶ Extended to modular-semilattices with applications to combinatorial optimizations [Hirai, Nakashima, 2020], [Hirai, Nakashima, 2018]
- ▶ MCCENUM *output-polynomial* for modular/median-semilattices [Hirai, Nakashima, 2018], [Kavvadias et al, 1995]
- ▶ Restricted case of dualization in lattices given by implicational bases, an *NP-complete problem* [Kavvadias et al, 1995], [Babin, Kuznetsov, 2017].

Quick Recap on Enumeration

- ▶ In enumeration, the size N of the output may be *exponential* in the size n .
- ▶ Let \mathcal{A} be an enumeration algorithm
 - ▶ Execution time bounded by $\text{poly}(n + N)$: \mathcal{A} runs in *output-polynomial* time,
 - ▶ Delay $\text{poly}(n)$ between two outputs: \mathcal{A} has *polynomial delay*
 - ▶ Delay $\text{poly}(n + i)$ between i -th and $i + 1$ -outputs: \mathcal{A} runs in *incremental-polynomial* time,
 - ▶ Execution time bounded by $2^{\text{polylog}(n+N)}$: \mathcal{A} runs in *output-quasipolynomial* time.



Execution time of \mathcal{A}



Connexion with co-atoms

- ▶ Using edges of G as new *keys*,
 - ▶ A *key* is a minimal subset K of X such that $K \rightarrow X$ holds.
- ▶ Maximal consistent closed sets are now *co-atoms*.

$$\Sigma = \left[\begin{array}{ll} 6 \rightarrow 2, & 16 \rightarrow 5, \\ 5 \rightarrow 2, & 15 \rightarrow 4, \\ 3 \rightarrow 1, & 24 \rightarrow 5, \\ 4 \rightarrow 1, & 16 \rightarrow X, \\ 56 \rightarrow X, & 35 \rightarrow X. \end{array} \right] \begin{array}{l} 2 \\ \bullet \end{array}$$



Connexion with co-atoms (bis)



- ▶ When adding edges of G as *keys*
- ▶ MCCENUM becomes a *restricted* instance of :

Problem - Co-atoms enumerations (CE)

- ▶ **Input:** A set of implications Σ over X .
 - ▶ **Output:** Co-atoms of \mathcal{F} .
-
- ▶ But, CE is untractable in output-poly time (unless $P = NP$) !
[Kavvadias et al, 1995]
 - ▶ So, what about our problem ?

Proof by picture

- ▶ Start from Σ over X (with induced \mathcal{F})
- ▶ Create fresh elements u, v , add $X \rightarrow uv$ to Σ
- ▶ The graph G (over $X \cup \{u, v\}$) has a unique edge: uv
- ▶ *maximal consistent closed sets* are duplications of \mathcal{F} 's *co-atoms*




Theorem [Nourine, V., 2021+]: The problem MCCENUM cannot be solved in output-polynomial time unless $P = NP$.

- ▶ Look carefully at the reduction of [Kavvadias et al, 1995]
- ▶ Slightly change the implication $X \rightarrow uv$
- ▶ Use the D-relation of [Freese et al, 1995] for lower bounded closure systems (*i.e. obtained by repeated duplications of lower-intervals*)

Corollary [Nourine, V., 2021+]: The problem MCCENUM cannot be solved in output-polynomial time unless $P = NP$, even when restricted to lower bounded closure systems.



An approach to tractable cases

- ▶ Co-atoms of are maximal independent sets of keys:
 - ▷ Add to Σ the implications $uv \rightarrow X$, for edges uv of G
 - ▷ Compute the keys \mathcal{K}
 - ▷ Find the maximal independent sets $\text{MIS}(\mathcal{K}) = \text{maxCC}(\Sigma, G)$.




How do keys behave ?

- ▶ In our case, a subset F must contain a key if:
 - ▶ it *contains* an edge of G ,
 - ▶ its *closure contains* an edge of G
- ▶ Thus, a *key* K has the form $K = A_u \cup A_v$ with A_u, A_v minimal generators of u, v
 - ▶ A_u *minimal generator* of u if it is a minimal subset of X such that $A_u \rightarrow u$.



Exponential Example


- ▶ $X = \{a_1, \dots, a_n, b_1, \dots, b_n, c_1, \dots, c_n, d_1, \dots, d_n, u, v\}$
- ▶ Σ with implications:
- ▶ G has a unique *edge* uv
- ▶ $\max\text{CC}(\Sigma, G)$ has *$2n$ solutions*: X minus a triple $\{u, a_i, b_i\}$ or $\{v, c_i, d_i\}$
- ▶ at least *2^{2n} keys* of *size $2n$* : all binary words on $\{a_i, b_i\}^n \times \{c_i, d_i\}^n$



Carathéodory number


- ▶ Carathéodory number $c(\mathcal{F})$: max. size of a minimal generator
- ▶ If bounded by a constant k :
 - ▶ Keys have size $\leq 2 \times k$
 - ▶ and \mathcal{K} has size $\text{poly}(|X|)$!
- ▶ $\text{MIS}(\mathcal{K}) = \max_{\text{CC}}(\Sigma, G)$ computable in incremental-poly time [Eiter et al., 1996]

Some closures with bounded $c(\mathcal{F})$




ideals of a poset

$$c(\mathcal{F}) = 1$$



convex subsets of a poset

$$c(\mathcal{F}) = 2$$



monophonic convexity of a chordal graph

$$c(\mathcal{F}) = 2$$



convex hull in \mathbb{R}^k

$$c(\mathcal{F}) = k + 1$$

The (atomistic) modular case

- ▶ MCCENUM easy to solve in distributive closure systems
- ▶ What about modularity ? Focus on the atomistic case:
 - ▷ *independence* of minimal generators [Grätzer, 2011]: subsets of A_u generates a boolean sublattice of \mathcal{F}
 - ▷ *biatomicity* [Bennett, 1987]: if $F \in \mathcal{F}$ and $F \cup x \rightarrow y$,

Theorem [Nourine, V., 2021+]: The problem MCCENUM can be solved in output-quasipolynomial time in atomistic modular closure systems.

MCCENUM : The big picture


output-poly

quasi-poly

\mathcal{K} exponential

untractable

unknown



Bool. = Boolean

Dist. = Distributive

At. = Atomistic

Mod. = Modular

Ac. = Acyclic

CG = Convex Geometry

Bd. = Bounded

LB = Lower Bounded

SD = Semidistributive

Conclusion

- ▶ Problem: given Σ and $G = (X, E)$, find maximal consistent (i.e independent) closed sets of \mathcal{F} w.r.t to G
- ▶ Results:
 - ✗ Not solvable in output-polynomial time unless $P = NP$,
 - ✓ Incremental-polynomial if the Carathéodory number is bounded,
 - ✓ Output-quasipolynomial in atomistic modular closure systems.
- ▶ Further research:
 - ▷ Tractability if the context is given as an input ?
 - ▷ Output-polynomial classes of closure systems generalizing distributivity ?

References

- ▶ **J-P. Barthélemy, J. Constantin**
Median graphs, parallelism and posets.
Discrete Mathematics, 111 :49-63, 1993.
- ▶ **M. Nielsen, G. Plotkin, G. Winskel**
Petri nets, event structures and domains, part I.
Theoretical Computer Science, 13 :85-108, 1981.
- ▶ **H. Hirai, S. Nakashima**
A compact representation for modular semilattices and its applications.
Order, 37 :479-507, 2020.
- ▶ **H. Hiroshi and T. Oki**
A compact representation for minimizers of k -submodular functions.
Journal of Combinatorial Optimization, 36 :709-741, 2018.
- ▶ **T. Eiter, G. Gottlob, K. Makino.**
New results on monotone dualization and generating hypergraph transversals.
SIAM Journal on Computing, 32 :514-537, 2003.
- ▶ **M. Fredman, L. Khachiyan.**
On the complexity of dualization of monotone disjunctive normal forms.
Journal of Algorithms, 21 :618-628, 1996.

References

- ▶ **D. Kavvadias, M. Sideri, E. Stavropoulos**
Generating all maximal models of a Boolean expression.
Information Processing Letters, 74 :157-162, 2000.
- ▶ **M. Babin, S. Kuznetsov**
Dualization in lattices given by ordered sets of irreducibles.
Theoretical Computer Science, 658 :316-326, 2017.
- ▶ **L. Nourine, S. Vilmin.**
Enumerating maximal consistent closed sets in closure systems
arXiv preprint arXiv:2102.04245, 2021.
- ▶ **M.K. Bennett**
Biatomic lattices.
Algebra Universalis, 24 :60-73, 1987.
- ▶ **G. Grätzer**
Lattice theory: foundation.
Springer Science & Business Media, 2011.
- ▶ **R. Freese, J. Ježek, J.B. Nation**
Free lattices.
American Mathematical Soc., 1995.